Article published on December 31st 2011 | Fitness

Ulcers can be very painful at times. Itâ€™s a disease caused by H. It is combined with the antibiotics amoxicillin and clarithromycin and within few days say 7 to 14 days it become worst. â€œPyloriâ€• is the root cause or majority causing factor of ulcers. When you know that you are suffering from ulcers and must contact the doctor at once? There are certain symptoms that you can notice yourself and it happen to you frequently. If you suffer any such symptoms you should immediately contact to doctor so that you are not suffering from any major disease and if you are it can be cure at once. Before talking further about it and its treatment you should know what ulcer actually is all about. In simple language one can say that itâ€™s a sore, a painful open wound which causes in stomach or upper part of small intestine.

The cause of ulcer can be simple that is stress, spicy food or alcohol. The most common sign of ulcers are stomach pain. If you continuously suffer from stomach pain then it would be advisable to concert a doctor as soon as possible otherwise it can grow larger or deeper. Other symptoms can be nausea, vomiting, weight loss. But if you know about your problem sooner than Nexium 40 mg is the best medicine you can go with. It helps you to get out of your problem of ulcers. For youngsters 40 mg dose is given daily for about 4- 8 weeks. For the treatment of H pylori, Nexium 40 mg is given for 10 days at one time in combination with other medicine known as amoxicillin and clarithromycin.

The storage of Nexium 40 mg is very important. You need to store these tablets carefully in a tight container and at room temperature carefully. It may happen that you suffer from some sort of problem after talking these medicines. The common side effects can be gas, headache, diarrhoea, dry mouth, abdominal pain, etc. Thus, it is advisable to concert the doctor before taking any medicine. It may happen you may suffer from serious problem after taking Nexium 40mg medicines like fever, stomach pain, tiredness, bleeding, yellowing of skin and eyes.

Always take this medicine with water. If a child suffers from ulcers you should ask the doctor to tell you the dose to give to the child as for different age groups the dose varies. It is always advisable to concert the doctor at once you suffer from any problem by taking Nexium 40 mg before stop the medicine. Keep the tablets out of reach of small children. Prevention is better than cure so it is advisable to wash your hand before and after every meal so that you keep yourself germ free and can avoid ulcers.

Article Source:
http://www.articleside.com/fitness-articles/know-what-ulcer-actually-is-all-about.htm - Article Side

Thakur7 - About Author:
Visit Trumedcanada.com which is a a Canadian Online Pharmacy and a Nexium 40mg

Article Keywords:
nexium generic, canadian online pharmacy, plavix 75mg, flomax, prilosec

You can find more free articles on Article Side. Sign up today and share your knowledge to the community! It is completely FREE!