Article Side

Enhancing Breast Symmetry through Breast Augmentation by Terry Bayer

Article published on March 20th 2012 | Health

Not all ladies are lucky to have perfect breasts; some have lopsided breasts that they are shy to even talk about. Most ladies who have asymmetrical breasts are those who experienced breastfeeding. Breastfeeding moms often give the same breast during feeding time causing uneven size. However, this must not be a cause for worry as there are various procedures that can help correct breast asymmetry. These procedures are simple and easily bearable for a lot of patients. Some doctors advise that women wait until age 20, when their breasts reached their full development, before they undergo any breast asymmetry procedure or any cosmetic procedure for the breast.

Some ladies have developed breasts but are not in proportion with their bodies. Others have small breasts but would rather have them fuller to allow them to wear clothes that look better with fuller breasts. Developments in cosmetic surgery now make it possible for full -breasted women to get breasts that are symmetrical; and for small-breasted women to have larger ones. The procedures are both quick and harmless.

For women with developed but uneven or lopsided breasts, a procedure to reduce breast asymmetry can be performed. This is may also include reduction of nipple size. The reconstructed breasts are reduced and lifted to look symmetrical.

Women with smaller breasts can go for implants to treat breast asymmetry. Implants can either be silicone or saline breast implants. Silicone implants are packed with silicone gel that seems like human fat. Some ladies prefer this type of implant over the other because it feels and appears more comfortable. Saline implants are filled with saline solution during the time of surgical treatment. Other patients say saline implants are not as soft and look unnatural. Patients can seek advice from their doctors on which is the safer and better option for them.

Patients who undergo any of the implant options need to put on special bras to help lessen breast swelling. These bras will provide the breasts enough support since inflammation and pain are expected for at least two days after the the treatment. Doctors recommend clients who have undergone breast augmentation Los Angeles clinics offer to use the special bra for several weeks and avoid doing physically demanding activities like lifting or pushing.

Cosmetic surgeons in Los Angeles are reputed to have more expertise in correcting breast asymmetry because of the wide client base they have. They can also conduct breast implant revision procedures if clients declare that they are not fully pleased with the breast procedure they had.

Article Source:
http://www.articleside.com/health-articles/enhancing-breast-symmetry-through-breast-augmentation.htm - Article Side

Terry Bayer - About Author:
For more details, search a breast asymmetry, a breast augmentation Los Angeles, and a breast implant revision in Google for related information.

Article Keywords:
breast asymmetry, breast augmentation Los Angeles, breast implant revision

You can find more free articles on Article Side. Sign up today and share your knowledge to the community! It is completely FREE!