Article published on May 1st 2012 | Health

A longstanding concern among a number of women is breast asymmetry-this causes a woman to have unequal breasts, or quite simply, one of the breasts is relatively smaller or larger compared with the other. Women of any age can suffer from this condition.

In most cases, breasts manifest themselves throughout a girlâ€™s adolescence, growing over time. Unfortunately, this is not what happens to 25 percent of the women globally who suffer from breast asymmetry. Research suggests that the normal weight of a breast must register at 16.9 ounces, and females having the condition will be ashamed about their light breasts.

Also, an irregular breast could be an underlying symptom of breast cancer. Based on the latest journal published on Breast Cancer Research, women that have asymmetrical breasts have a fifty percent risk of getting cancer of the breast. Thatâ€™s why itâ€™s essential for females having this probler to have doctors look at their asymmetrical breasts to know if there are actually problems they need to be aware of. Researchers, however, noted that it's natural for women to have partially different breast shapes or sizes; whatâ€™s scary is the case of close in size breasts eventually getting asymmetrical.

Genetics

Medical professionals report that genetics or haphazard growth patterns in the body are a contributing component why several women acquire breast asymmetry. The human breast forms due to the female hormone estrogen, expanding two years right after the girl experiences her very first menstrual period. Over the next 2 to 4 years, the breast is going to continue developing right until it attains maturity. This is not the truth for women who have abnormal breast growths.

Cancer of the Breast

Based on health news reports, an irregular breast can be an indicator of imminent illnesses, such as breast cancer. Medical professionals often recommend breast augmentation Los Angeles health centers offer to address the results of breast asymmetry, but self-breast examinations and routine mammogram screening should be done appropriately.

Pregnancy and Sagging

There are tons of hormonal changes happening in a pregnant womanâ€™s body. Her breasts become extra tender throughout the entire pregnancy phase. She might discover that her left breast is larger than the other as a result of the production of milk, or lactation. Meanwhile, ptosis or sagging occurs when a woman starts to age over time. Adult women who went through breast augmentation procedures may also find their chests drooping. This can be fixed by having breast implant revision executed on both breasts to improve their overall look and feel. Go to wdxcyber.com/nbreast3.htm for more information on the major triggers of breast asymmetry.

Article Source:
http://www.articleside.com/health-articles/fundamental-information-regarding-asymmetrical-breasts.htm - Article Side

Terry Bayer - About Author:
For more details, search a breast implant revision, a breast augmentation Los Angeles and a breast asymmetry in Google for more related information.

Article Keywords:
breast asymmetry, breast augmentation los angeles, breast implant revision

You can find more free articles on Article Side. Sign up today and share your knowledge to the community! It is completely FREE!